

Isostaticity in two dimensional pile of rigid disks

Akihiro KASAHARA and Hiizu NAKANISHI*

Department of Physics, Kyushu University 33, Fukuoka 812-8581

We study the static structure of piles made of polydisperse disks in the rigid limit with and without friction using molecular dynamic simulations for various elasticities of the disks and pile preparation procedures. The coordination numbers are calculated to examine the isostaticity of the pile structure. For the frictionless pile, it is demonstrated that the coordination number converges to 4 in the rigid limit, which implies that the structure of rigid disk pile is isostatic. On the other hand, for the frictional case with the infinite friction constant, the coordination number depends on the preparation procedure of the pile, but we find that the structure becomes very close to isostatic with the coordination number close to 3 in the rigid limit when the pile is formed through the process that tends to make a pile of random configuration.

KEYWORDS: static structure of granular pile, isostaticity, marginal rigidity, stress propagation

Isostaticity is a peculiar property that a rigid granular pile is expected to have.¹⁾ Consider first a pile of frictionless deformable grains under gravity. The pile is assumed to be stable against small enough external perturbation. Each grain in the pile should be deformed and the forces acting through the contacts between the grains are elastic force, that is related to the grain deformation through elasticity. Imagine that the grains in the pile are being hardened gradually. As the grains become harder, their deformation diminishes and some of the contacts disappear. In the rigid limit, no deformation is allowed and the forces acting between the grains should be determined only by the structure of contact network of the pile without any information of deformation.

Based on such observation, it has been conjectured¹⁾ that the contact network in a static pile of non-cohesive rigid grains without friction should have a property that the network has just enough contacts to be rigid in the sense that removal of any contact makes the structure flexible. This property is called *isostaticity*^{1,2)} or *marginal rigidity*.^{3,4)} the structure should be unstable against an infinitesimal small external force if there are less contacts than this, while it should be overconstrained with internal stress if there are more contacts.

It is not obvious, however, whether this holds true or not for the pile that is formed through deposition of grains that are rigid from the beginning. It is even less obvious for the case that the friction between the grains comes into the problem.

A simple way to check whether a pile is isostatic is just to count the average coordination number z of grains in the pile. Since all the contact forces in the isostatic pile have to be determined without redundant constraint, the number of independent components of the forces should coincide with the number of the conditions for the force balance and the torque balance for each grain. This leads to $z = 2d$ for a pile of frictionless spheres, $z = d(d+1)$ for a pile of frictionless non-spherical grains, and $z = d + 1$ for a frictional pile of both spherical and non-spherical grains in d dimensions.

Isostaticity for the piles deposited from a certain initial configuration has been examined by numerical simulations for three dimensional monodisperse spherical systems with various elastic constants,⁵⁾ and it was shown that the piles of frictionless sphere becomes isostatic in the hard sphere limit, while the pile of frictional spheres are more dependent on the way how the piles are prepared, but are not likely to be isostatic.

On the other hand, Blumenfeld et al. performed simple tabletop experiments for two dimensional non-spherical grains of cardboard and concluded that the piles of two dimensional grains become isostatic in the limit that the number density of a initial configuration and that of a final stable configuration coincides.

In this report, using numerical simulations for two dimensional system, we examine the piles of two dimensional polydisperse disk system for several ways of preparation procedures.

We performed molecular dynamics simulations on the system that consists of two dimensional disks with linear elasticity and damping. The system is polydisperse with the uniform distribution in the disk diameter with the range $[0.9, 1]$ of the maximum diameter σ_0 , and their masses are assumed to be proportional to their areas:




Fig. 1. Two initial configurations: the triangular lattice with polydisperse disks (left) and the random configuration with the number density 0.6 (right).

*E-mail address: naka4scp@mbox.nc.kyushu-u.ac.jp




Fig. 2. The coordination number z for various elastic constant k_n for frictionless pile (a) and frictional pile (b). The marks represent the pile preparation procedure: the Newtonian equation with the triangular lattice initial configuration (+), the Newtonian equation with the random initial configuration (x), the viscous equation with the triangular lattice initial configuration (*), the viscous equation with the random initial configuration (□). Each mark represents average over six to twelve realizations.

the mass of the disk with the diameter σ_0 is denoted by m_0 . The bottom of the system is made rough by attaching the disks with the interval σ_0 , and we employ the periodic boundary condition in the horizontal direction. The number of the disks N in the system is 400, and the horizontal length of the system is $20\sigma_0$, thus we have approximately 20 layers of disks on average.

Piles are formed by letting the system run under the gravitational acceleration g from initial configurations until all the disks stop moving.

The disk at the position $\mathbf{x}(t)$ with mass m follows the Newtonian equation

$$m\ddot{\mathbf{x}}(t) = \mathbf{F}(t), \quad (1)$$

where the force $\mathbf{F}(t)$ consists of the gravitation and the contact force from the neighboring disks in contact. The two particles i and j at $\mathbf{x}_i(t)$ and $\mathbf{x}_j(t)$ with radii r_i and




Fig. 3. The coordination numbers z for various elastic constant k_n in the log-log scale frictionless pile (a) and frictional pile (b). The same data are plotted using the same marks with those in Fig.2

r_j , respectively, are in contact when the overlap δ_{ij} given by

$$\delta_{ij} = r_i + r_j - |\mathbf{x}_i - \mathbf{x}_j|, \quad (2)$$

is positive. Then the particle j exerts the force $\mathbf{F}_{ij}$ on the particle i

$$\mathbf{F}_{ij} = \mathbf{F}_{ij}^n + \mathbf{F}_{ij}^t, \quad (3)$$

where $\mathbf{F}_{ij}^n$ and $\mathbf{F}_{ij}^t$ are the normal and tangential components of the force;

$$\mathbf{F}_{ij}^n = k_n \delta_{ij} \hat{\mathbf{n}}_{ij} - \gamma_n \mathbf{v}_{ij}^n, \quad (4)$$

$$\mathbf{F}_{ij}^t = -k_t \Delta s_{ij} \hat{\mathbf{t}}_{ij} - \gamma_t \mathbf{v}_{ij}^t. \quad (5)$$

Here, $\hat{\mathbf{n}}_{ij}$ and $\hat{\mathbf{t}}_{ij}$ are the normal and tangential unit vectors, respectively; Δs_{ij} is the tangential displacement of the contact points after the contact, $\mathbf{v}_{ij}^n$ ($\mathbf{v}_{ij}^t$) is relative normal (tangential) velocity, k_n (k_t) is the normal (tangential) elastic constant, and γ_n (γ_t) is the normal (tangential) damping constant. Note that we assume no

Eq. of motion	Initial config.	Frictionless		Frictional	
		z_∞	α	z_∞	α
Newtonian	Triangular lattice	3.97	0.68	3.15	0.49
	Random config.	3.98	0.65	3.09	0.47
Viscous	Triangular lattice	3.97	0.63	3.06	0.60
	Random config.	3.97	0.64	3.04	0.46

Table I. The limiting coordination numbers z_∞ and the exponents α for various preparation procedures.

threshold for the disks to slip during the contact, which corresponds to the case with the infinite friction coefficient. In the case of frictionless disk, we simply set $F_{ij}^t = 0$.

In the actual simulations, we use $k_t = 0.2k_n$ and $\gamma_n = 2\sqrt{k_n}[\sqrt{m_0}]$ for the frictionless case, and $k_t = 0.2k_n$ and $\gamma_n = \gamma_t = 2\sqrt{k_n}[\sqrt{m_0}]$ for the frictional case.

As for the initial configurations, we try the triangular lattice and the random configuration (Fig.1); the triangular lattice with the lattice constant σ_0 is not a regular lattice because the disks located at the lattice points are polydisperse. The random configurations are prepared by putting disks at random with the area fraction around 0.6.

Simulations to form piles start from these configurations with zero particle velocity and stop when the kinetic energy of each disks becomes negligibly small, namely smaller than $10^{-15}[m_0 g \sigma_0]$.

Fig.2(a) shows the k_n -dependence of the coordination number z for the frictionless disks. It can be seen that the preparation dependence is very small and z converges to the number very close to 4 in the large k_n limit for both of the initial configurations. The k_n -dependence is well represented by the power law

$$z - z_\infty \propto k_n^{-\alpha} \quad (6)$$

as is shown in Fig.3(a).

As for the case of frictional disks, the results are shown in Fig. 2(b). There are two things to be noted in comparison with the frictionless case: (i) the discrepancy between the two initial conditions shown by + and $\times$ is large, (ii) the limiting values of the coordination number is substantially different from 3, or the value for the isostatic structure of the frictional grain in two dimensions.

The limiting values of the coordination numbers are $z_\infty = 3.15$ for the triangular lattice initial condition and $z_\infty = 3.09$ for the random initial condition (Table I).

These features indicate that the piles of frictional disks are more dependent on the way they are formed than those of the frictionless piles. The fact that z_∞ for the random initial conditions is closer to 3 suggests that the piles would tend to be isostatic when they are formed through the process that retains a random structure.

This consideration prompts us to employ the viscous equation

$$\gamma \dot{\mathbf{x}}(t) = \mathbf{F}(t) \quad (7)$$

for the time development during the deposition process in stead of the Newtonian equation, in order to see if it makes the structure closer to the isostaticity. The viscous equation tends to produce random configurations in equilibrium because the disks stop as soon as the force balance is achieved without the inertia effect. In the simulation with the viscous equation, we take $\gamma = 5[m_0\sqrt{g/\sigma_0}]$ with $\gamma_n = \gamma_t = 0$ in eqs.(4) and (5).

The data for the piles by the viscous equation are also shown in Fig.3 and Table I. One can see that the limiting values for the viscous equation are actually closer to 3 than those for the Newtonian equation.

In summary, in this report we have shown the followings; The piles of frictionless disks become isostatic when the disks are very hard and they are not sensitive to the preparation procedure very much, which is consistent with the conjecture that the pile of rigid grains is isostatic. On the other hand, for the piles of frictional disks with the infinite friction, the structure depends on the preparation process. If the pile is formed from the triangular lattice with the inertia, the pile structure seems to be distinctively different from the isostatic one even in the rigid limit, as has been found in the previous work on the three dimensional system.⁵⁾ We found, however, that the pile of frictional disks becomes very close to isostatic in the rigid grain limit when we employ the deposition process that tends to produce random structure.

The isostaticity of granular pile should play an crucial role in the stress propagation of granular system. In the case of frictionless pile, it has been shown that the isostatic structure has some peculiar features: (i) unidirectional nature of stress propagation,⁶⁾ (ii) correspondence between the force-force response and the displacement-displacement response,²⁾ (iii) unstable response that grows as the distance from the point of an external perturbation.¹⁾ The effects of friction on the mechanical properties of the isostatic structure are now under investigation.

- 1) C.F. Moukarzel, Phys. Rev. Lett. **81** (1998) 1634.
- 2) C.F. Moukarzel, Granular Matter, **3** (2001) 41.
- 3) R.C. Ball and R. Blumenfeld, Phys. Rev. Lett. **88** (2002) 115505.
- 4) R.C. Ball and R. Blumenfeld, Phil. Trans. R. Soc. Lond. A **361** (2003) 731.
- 5) L.E. Silbert, D.Ertas, G.S. Grest, T.C. Halsey, and D. Levine, Phys. Rev. E **65** (2002) 031304.
- 6) A.V. Tkachenko and T.A. Witten, Phys. Rev. E **60** (1999) 687.